# *Australia New Zealand Food Standards Code* – Amendment No. 86 – 2006

## *Food Standards Australia New Zealand Act 1991*

### Preamble

The variations set forth in the Schedule below are variations to Standards in the *Australia New Zealand Food Standards Code* published by the National Health and Medical Research Council in the *Commonwealth of Australia Gazette*, No. P 27, on 27 August 1987, which have been varied from time to time.

These variations are published pursuant to section 23A of the *Food Standards Australia New Zealand Act 1991*.

### Citation

These variations may be collectively known as the *Australia New Zealand Food Standards Code* – Amendment No. 86 – 2006.

### Commencement

These variations commence on Gazettal.

### SCHEDULE

**[1]** ***Standard 1.3.3*** *is varied by –*

[1.1] i*nserting into the* Table to clause 17 –

| | |
|---|---|
| Lipase, triacylglycerol<br>EC [3.1.1.3] | *Mucor javanicus* |

[1.2] *inserting into the* Editorial note *to clause 17 –*

| |
|---|
| *Mucor javanicus* is also known as *Mucor circinelloides* f. *circinelloides*. |

**[2]** ***Standard 1.4.2*** *is varied by –*

[2.1] *omitting from* Schedule 1 *all entries for the following chemicals –*

Alloxydim
Alloxydim sodium
Diclobutrazol
Diofenolan
Diphenamid
Methazole
Promecarb



Federal Register of Legislative Instruments F2006L01578

[2.2] *omitting from* Schedule 1 *the chemical and chemical residue definition appearing in* Column 1 *of the Table to this sub-item, substituting the chemical and chemical residue definition in appearing in* Column 2 –

| COLUMN 1 | COLUMN 2 |
|---|---|
| **TYLOSIN**<br>TYLOSIN | **TYLOSIN**<br>TYLOSIN A |

[2.3] *inserting in* Schedule 1 –

| **CLOTHIANIDIN**<br>*COMMODITIES OF PLANT ORIGIN*:<br>*COMMODITIES OF ANIMAL ORIGIN*: SUM OF CLOTHIANIDIN, 2-CHLOROTHIAZOL-5-YLMETHYLGUANIDINE, 2-CHLOROTHIAZOL-5-YLMETHYLUREA, AND THE PYRUVATE DERIVATIVE OF N'-(2-CHLOROTHIAZOL-5-YLMETHYL)-N'-METHYLGUANIDINE, EXPRESSED AS CLOTHIANIDIN | |
|---|---|
| COTTON SEED | T*0.02 |
| EDIBLE OFFAL (MAMMALIAN) | T*0.02 |
| MEAT (MAMMALIAN) (IN THE FAT) | T*0.02 |
| MILKS | T*0.01 |
| **FLUMICLORAC PENTYL**<br>FLUMICLORAC PENTYL | |
| COTTON SEED | T0.1 |
| EDIBLE OFFAL (MAMMALIAN) | T*0.01 |
| EGGS | T*0.01 |
| MEAT (MAMMALIAN) | T*0.01 |
| MILKS | T*0.01 |
| POULTRY, EDIBLE OFFAL OF | T*0.01 |
| POULTRY MEAT | T*0.01 |
| **FORCHLORFENURON**<br>FORCHLORFENURON | |
| GRAPES | T*0.01 |
| **METHYL ISOTHIOCYANATE**<br>METHYL ISOTHIOCYANATE | |
| BARLEY | T0.1 |
| RAPE SEED | T0.1 |
| WHEAT | T0.1 |
| **ROBENIDINE**<br>ROBENIDINE | |
| POULTRY, EDIBLE OFFAL OF | *0.1 |
| POULTRY MEAT | *0.1 |

[2.4] *omitting from* Schedule 1 *the foods and associated MRLs for each of the following chemicals –*

| **BENFLURALIN**<br>BENFLURALIN | |
|---|---|
| EDIBLE OFFAL (MAMMALIAN) | T*0.01 |
| MEAT (MAMMALIAN) | T*0.01 |
| MILKS | T*0.01 |
| **CARBENDAZIM**<br>SUM OF CARBENDAZIM AND 2-AMINOBENZIMIDAZOLE, EXPRESSED AS CARBENDAZIM | |
| BROAD BEANS (DRY) | T0.5 |
| CHICK-PEA (DRY) | T0.5 |
| LENTILS (DRY) | T0.5 |
| TREE NUTS | T0.1 |
| **CHLORPYRIFOS-METHYL**<br>CHLORPYRIFOS-METHYL | |
| COTTON SEED OIL, CRUDE | *0.01 |
| **CYPROCONAZOLE**<br>CYPROCONAZOLE, SUM OF ISOMERS | |
| BANANA | T0.5 |
| GRAPES | T0.5 |
| **DIFENOCONAZOLE**<br>DIFENOCONAZOLE | |
| CEREAL GRAINS | T*0.01 |
| **DIMETHOMORPH**<br>SUM OF E AND Z ISOMERS OF DIMETHOMORPH | |
| CHARD (SILVER BEET) | T2 |
| LETTUCE, LEAF | T2 |
| **DIQUAT**<br>DIQUAT CATION | |
| COTTON SEED | 1 |
| COTTON SEED OIL, CRUDE | 0.1 |
| LENTIL (DRY) | T0.5 |
| LUPIN (DRY) | 0.5 |
| POPPY SEED | 5 |
| RAPE SEED | 2 |
| RAPE SEED OIL, CRUDE | 0.1 |
| SESAME SEED | 5 |
| SESAME SEED OIL, CRUDE | 0.1 |
| SOYA BEAN (DRY) | 1 |

| | |
|---|---|
| SUNFLOWER SEED | 1 |
| SUNFLOWER SEED OIL, CRUDE | 1 |

| **DITHIOCARBAMATES**<br>TOTAL DITHIOCARBAMATES, DETERMINED AS CARBON DISULPHIDE EVOLVED DURING ACID DIGESTION AND EXPRESSED AS MILLIGRAMS OF CARBON DISULPHIDE PER KILOGRAM OF FOOD | |
|---|---|
| BROAD BEANS (DRY) (FAVA BEAN) | 0.5 |
| CHICK-PEA (DRY) | 0.5 |
| HERBS [EXCEPT PARSLEY] | T5 |
| LENTIL (DRY) | 0.5 |
| PEAS (DRY) | T0.5 |

| **ETHOPROPHOS**<br>ETHOPROPHOS | |
|---|---|
| GRAPES | T*0.01 |

| **FENOXYCARB**<br>FENOXYCARB | |
|---|---|
| GRAPES | T2 |
| STONE FRUITS | T0.5 |

| **FLUVALINATE**<br>FLUVALINATE, SUM OF ISOMERS | |
|---|---|
| CHERRIES | T*0.05 |
| NECTARINE | 0.1 |
| PEACH | T0.1 |
| PLUMS (INCLUDING PRUNES) | T0.1 |

| **LINURON**<br>SUM OF LINURON PLUS 3,4-DICHLOROANILINE, EXPRESSED AS LINURON | |
|---|---|
| VEGETABLES [EXCEPT LEEK] | *0.05 |

| **METALAXYL**<br>METALAXYL | |
|---|---|
| CEREAL GRAINS | T*0.05 |
| HERBS | T0.3 |
| MEAT (MAMMALIAN) (IN THE FAT) | *0.05 |

| **METHOMYL**<br>SUM OF METHOMYL AND METHYL HYDROXYTHIOACETIMIDATE ('METHOMYL OXIME'), EXPRESSED AS METHOMYL<br>*SEE ALSO* THIODICARB | |
|---|---|
| LEAFY VEGETABLES | 1 |

| **PHOSPHOROUS ACID**<br>PHOSPHOROUS ACID | |
|---|---|
| APPLE | 50 |
| AVOCADO | 100 |
| CHERVIL | T5 |
| CHESTNUTS | T500 |
| CUCURBITS | 25 |
| DURIAN | T100 |
| GRAPE LEAVES | 300 |
| GRAPES | 50 |
| PEACH | 100 |
| PINEAPPLE | 50 |
| PISTACHIO NUT | T1000 |
| RASPBERRIES | T50 |
| RUCOLA (ROCKET) | T5 |
| STRAWBERRY | T50 |
| TURMERIC, ROOT | T5 |
| WALNUTS | T50 |

| **PROCYMIDONE**<br>PROCYMIDONE | |
|---|---|
| CARROT | T1 |
| GRAPES | 2 |
| LETTUCE, HEAD | 2 |
| LETTUCE, LEAF | 2 |
| STRAWBERRY | 5 |
| TOMATO | 2 |

| **PROPACHLOR**<br>PROPACHLOR | |
|---|---|
| CEREAL GRAINS | *0.05 |

| **SETHOXYDIM**<br>SUM OF SETHOXYDIM AND METABOLITES CONTAINING THE 5-(2-ETHYLTHIOPROPYL)CYCLOHEXENE-3-ONE AND 5-HYDROXYCYCLOHEXENE-3-ONE MOIETIES AND THEIR SULFOXIDES AND SULFONES, EXPRESSED AS SETHOXYDIM | |
|---|---|
| PEANUT OIL, CRUDE | 2 |

| **TRICHLORFON**<br>TRICHLORFON | |
|---|---|
| PEPPERS, SWEET | T0.5 |

[2.5] *inserting in alphabetical order in* Schedule 1, *the foods and associated MRLs for each of the following chemicals –*

| **ABAMECTIN**<br>SUM OF AVERMECTIN B1A, AVERMECTIN B1B AND (Z)-8,9 AVERMECTIN B1A, AND (Z)-8,9 AVERMECTIN B1B | |
|---|---|
| CUCUMBER | 0.02 |
| PEAS | T0.2 |
| SQUASH, SUMMER | 0.02 |

| **AZOXYSTROBIN**<br>AZOXYSTROBIN | |
|---|---|
| BEANS [EXCEPT BROAD AND SOYA BEAN] | T3 |

| | |
|---|---|
| LETTUCE, HEAD | T3 |
| LETTUCE, LEAF | T3 |
| OLIVES | T2 |

| **BIFENTHRIN**<br>BIFENTHRIN | |
|---|---|
| PEAS (PODS AND SUCCULENT, IMMATURE SEEDS) | T*0.01 |

| **BOSCALID**<br>*COMMODITIES OF PLANT ORIGIN:* BOSCALID<br>*COMMODITIES OF ANIMAL ORIGIN:* SUM OF BOSCALID, 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE AND GLUCURONIDE CONJUGATE OF 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE, EXPRESSED AS BOSCALID EQUIVALENTS | |
|---|---|
| BEANS [EXCEPT BROAD BEAN AND SOYA BEAN] | T3 |
| BRASSICA (COLE OR CABBAGE) VEGETABLES, HEAD CABBAGES, FLOWERHEAD BRASSICAS | T2 |
| BRASSICA LEAFY VEGETABLES | T10 |
| LETTUCE, HEAD | T2 |
| LETTUCE, LEAF | T2 |
| ONION, BULB | T1.0 |
| STRAWBERRY | T5 |

| **BUPIRIMATE**<br>BUPIRIMATE | |
|---|---|
| PEPPERS | T1 |

| **CARBENDAZIM**<br>SUM OF CARBENDAZIM AND 2-AMINOBENZIMIDAZOLE, EXPRESSED AS CARBENDAZIM | |
|---|---|
| MACADAMIA NUTS | 0.1 |
| PISTACHIO NUT | T0.1 |
| PULSES | 0.5 |

| **CHLORMEQUAT**<br>CHLORMEQUAT CATION | |
|---|---|
| BARLEY | T2 |
| EDIBLE OFFAL (MAMMALIAN) | 0.5 |
| EGGS | 0.1 |
| MEAT (MAMMALIAN) | 0.2 |
| POULTRY, EDIBLE OFFAL OF | 0.1 |
| POULTRY MEAT | *0.05 |

| **CHLORPYRIFOS**<br>CHLORPYRIFOS | |
|---|---|
| STAR APPLE | T*0.05 |

| **CYHALOTHRIN**<br>CYHALOTHRIN, SUM OF ISOMERS | |
|---|---|
| CUCUMBER | T0.05 |

| **CYPERMETHRIN**<br>CYPERMETHRIN, SUM OF ISOMERS | |
|---|---|
| CORIANDER (LEAVES, STEM, ROOTS) | T1 |
| CORIANDER, SEED | T1 |
| PARSLEY | T1 |

| **DIFENOCONAZOLE**<br>DIFENOCONAZOLE | |
|---|---|
| BARLEY | *0.01 |
| WHEAT | *0.01 |

| **DIMETHOMORPH**<br>SUM OF E AND Z ISOMERS OF DIMETHOMORPH | |
|---|---|
| LEAFY VEGETABLES [EXCEPT LETTUCE HEAD] | T2 |

| **DIQUAT**<br>DIQUAT CATION | |
|---|---|
| OILSEED [EXCEPT LINSEED] | 5 |
| PULSES | 1 |
| VEGETABLE OILS, CRUDE | 1 |

| **DITHIOCARBAMATES**<br>TOTAL DITHIOCARBAMATES, DETERMINED AS CARBON DISULPHIDE EVOLVED DURING ACID DIGESTION AND EXPRESSED AS MILLIGRAMS OF CARBON DISULPHIDE PER KILOGRAM OF FOOD | |
|---|---|
| PULSES | 0.5 |
| RADISH | T1 |
| SWEDE | T1 |
| TURNIP, GARDEN | T1 |

| **EPOXICONAZOLE**<br>EPOXICONAZOLE | |
|---|---|
| BARLEY | T0.5 |
| EGGS | T*0.01 |
| POULTRY, EDIBLE OFFAL OF | T0.02 |
| POULTRY MEAT (IN THE FAT) | T0.05 |
| WHEAT | T0.5 |
| WHEAT BRAN, UNPROCESSED | T3 |
| WHEAT GERM | T2 |

| **ETHEPHON**<br>ETHEPHON | |
|---|---|
| WALNUTS | T0.5 |

| **FIPRONIL**<br>SUM OF FIPRONIL, THE SULPHENYL METABOLITE (5-AMINO-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-4-[(TRIFLUOROMETHYL) SULPHENYL]-1H-PYRAZOLE-3-CARBONITRILE), THE SULPHONYL METABOLITE (5-AMINO-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-4-[(TRIFLUOROMETHYL)SULPHONYL]-1H-PYRAZOLE-3-CARBONITRILE), AND THE TRIFLUOROMETHYL METABOLITE (5-AMINO-4-TRIFLUOROMETHYL-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-1H-PYRAZOLE-3-CARBONITRILE) | |
|---|---|
| HONEY | T0.05 |
| **FLUVALINATE**<br>FLUVALINATE, SUM OF ISOMERS | |
| STONE FRUITS | 0.05 |
| **GLYPHOSATE**<br>SUM OF GLYPHOSATE AND AMINOMETHYLPHOSPHONIC ACID (AMPA) METABOLITE, EXPRESSED AS GLYPHOSATE | |
| SORGHUM | T10 |
| **IMAZALIL**<br>IMAZALIL | |
| MELONS [EXCEPT WATERMELON] | 10 |
| **IPRODIONE**<br>IPRODIONE | |
| ONION, BULB | T0.2 |
| **LINURON**<br>SUM OF LINURON PLUS 3,4-DICHLOROANILINE, EXPRESSED AS LINURON | |
| CELERY | *0.05 |
| VEGETABLES [EXCEPT CELERY AND LEEK] | *0.05 |
| **MALEIC HYDRAZIDE**<br>SUM OF FREE AND CONJUGATED MALEIC HYDRAZIDE, EXPRESSED AS MALEIC HYDRAZIDE | |
| CARROT | T40 |
| **MELOXICAM**<br>MELOXICAM | |
| PIG FAT/SKIN | 0.1 |
| PIG KIDNEY | *0.01 |
| PIG LIVER | *0.01 |
| PIG MEAT | 0.02 |
| **METALAXYL**<br>METALAXYL | |
| BARLEY | *0.01 |
| EGGS | *0.05 |
| MEAT (MAMMALIAN) | *0.05 |
| POULTRY, EDIBLE OFFAL OF | *0.05 |
| POULTRY MEAT | *0.05 |
| WHEAT | *0.01 |
| **METHOMYL**<br>SUM OF METHOMYL AND METHYL HYDROXYTHIOACETIMIDATE ('METHOMYL OXIME'), EXPRESSED AS METHOMYL<br>*SEE ALSO* THIODICARB | |
| CHARD | T2 |
| LEAFY VEGETABLES [EXCEPT CHARD] | 1 |
| **METRIBUZIN**<br>METRIBUZIN | |
| SUGAR CANE | 0.1 |
| **PHENMEDIPHAM**<br>PHENMEDIPHAM | |
| LETTUCE, HEAD | T0.2 |
| LETTUCE, LEAF | T0.2 |
| **PHOSPHOROUS ACID**<br>PHOSPHOROUS ACID | |
| ASSORTED TROPICAL AND SUBTROPICAL FRUITS – INEDIBLE PEEL | T100 |
| BERRIES AND OTHER SMALL FRUITS | T50 |
| BULB VEGETABLES | T10 |
| KAFFIR LIME LEAVES | T5 |
| LEAFY VEGETABLES | T100 |
| LEMON GRASS | T5 |
| LEMON VERBENA | T5 |
| PEAS, SHELLED | T100 |
| POPPY SEED | 1 |
| ROOT AND TUBER VEGETABLES | T100 |
| TREE NUTS | T1000 |
| **PICOLINAFEN**<br>*COMMODITIES OF PLANT ORIGIN*: PICOLINAFEN<br>*COMMODITIES OF ANIMAL ORIGIN*: SUM OF PICOLINAFEN AND 6-[3-TRIFLUOROMETHYL PHENOXY]-2-PYRIDINE CARBOXYLIC ACID | |
| EGGS | *0.01 |
| POULTRY, EDIBLE OFFAL OF | *0.02 |
| POULTRY MEAT (IN THE FAT) | *0.02 |
| **PROCYMIDONE**<br>PROCYMIDONE | |
| PEPPERS | T2 |
| ROOT AND TUBER VEGETABLES [EXCEPT POTATO] | T1 |
| WINE GRAPES | T2 |
| **PROPACHLOR**<br>PROPACHLOR | |
| CEREAL GRAINS [EXCEPT SORGHUM] | 0.05 |

| | |
|---|---|
| EDIBLE OFFAL (MAMMALIAN) | 0.1 |
| EGGS | *0.02 |
| MEAT (MAMMALIAN) (IN THE FAT) | *0.02 |
| MILKS | *0.02 |
| POULTRY, EDIBLE OFFAL OF | *0.02 |
| POULTRY MEAT (IN THE FAT) | *0.02 |
| SORGHUM | 0.2 |
| SWEET CORN (CORN-ON-THE-COB) | 0.05 |
| TURNIP, GARDEN | *0.02 |
| **PYMETROZINE** PYMETROZINE | |
| BEETROOT | *0.02 |
| **SETHOXYDIM** SUM OF SETHOXYDIM AND METABOLITES CONTAINING THE 5-(2-ETHYLTHIOPROPYL)CYCLOHEXENE-3-ONE AND 5-HYDROXYCYCLOHEXENE-3-ONE MOIETIES AND THEIR SULFOXIDES AND SULFONES, EXPRESSED AS SETHOXYDIM | |
| BARLEY | *0.1 |
| LINSEED | 0.5 |
| SPRING ONION | T0.5 |
| **SPINOSAD** SUM OF SPINOSYN A AND SPINOSYN D | |
| SAFFLOWER SEED | T*0.01 |

| | |
|---|---|
| **TOLCLOFOS-METHYL** TOLCLOFOS-METHYL | |
| BEETROOT | T0.5 |
| **TOLTRAZURIL** SUM OF TOLTRAZURIL, ITS SULFOXIDE AND SULFONE, EXPRESSED AS TOLTRAZURIL | |
| EGGS | T*0.05 |
| **TOLYLFLUANID** TOLYLFLUANID | |
| BERRIES AND OTHER SMALL FRUITS [EXCEPT GRAPES AND STRAWBERRY] | T15 |
| DRIED GRAPES | T0.2 |
| GRAPES | T*0.05 |
| **TRICHLORFON** TRICHLORFON | |
| GOAT, EDIBLE OFFAL OF | 0.1 |
| GOAT MEAT | 0.1 |
| PEPPERS | 0.2 |
| **TRICLOPYR** TRICLOPYR | |
| CITRUS FRUITS | T0.1 |

[2.6] *omitting from* Schedule 1*, under the entries for the following chemicals, the maximum residue limit for the food, substituting –*

| | |
|---|---|
| **ABAMECTIN** SUM OF AVERMECTIN B1A, AVERMECTIN B1B AND (Z)-8,9 AVERMECTIN B1A, AND (Z)-8,9 AVERMECTIN B1B | |
| EGGPLANT | 0.02 |
| **AZOXYSTROBIN** AZOXYSTROBIN | |
| COTTON SEED | *0.01 |
| **CHLORMEQUAT** CHLORMEQUAT CATION | |
| MILKS | 0.5 |
| **CYPERMETHRIN** CYPERMETHRIN, SUM OF ISOMERS | |
| LINOLA OIL, EDIBLE | 0.1 |
| LINOLA SEED | 0.1 |

| | |
|---|---|
| **DITHIOCARBAMATES** TOTAL DITHIOCARBAMATES, DETERMINED AS CARBON DISULPHIDE EVOLVED DURING ACID DIGESTION AND EXPRESSED AS MILLIGRAMS OF CARBON DISULPHIDE PER KILOGRAM OF FOOD | |
| ALMONDS | 3 |
| BEETROOT | 1 |
| CITRUS FRUITS | 0.2 |
| COTTON SEED | 10 |
| CUSTARD APPLE | 5 |
| POME FRUITS | 3 |
| POTATO | 1 |
| STRAWBERRY | 3 |
| **DODINE** DODINE | |
| STONE FRUITS | *0.05 |
| **EPOXICONAZOLE** EPOXICONAZOLE | |
| EDIBLE OFFAL (MAMMALIAN) | T0.05 |
| MILKS | T0.01 |

| **FLUAZIFOP-BUTYL**<br>FLUAZIFOP-BUTYL | |
|---|---|
| GINGER, ROOT | 0.05 |

| **FLUDIOXONIL**<br>*COMMODITIES OF ANIMAL ORIGIN:* SUM OF FLUDIOXONIL AND OXIDISABLE METABOLITES, EXPRESSED AS FLUDIOXONIL<br>*COMMODITIES OF PLANT ORIGIN:* FLUDIOXONIL | |
|---|---|
| COTTON SEED | *0.05 |
| RAPE SEED | *0.01 |

| **FLUVALINATE**<br>FLUVALINATE, SUM OF ISOMERS | |
|---|---|
| COTTON SEED | 0.1 |
| TABLE GRAPES | 0.05 |

| **GLYPHOSATE**<br>SUM OF GLYPHOSATE AND AMINOMETHYLPHOSPHONIC ACID (AMPA) METABOLITE, EXPRESSED AS GLYPHOSATE | |
|---|---|
| CEREAL GRAINS [EXCEPT AS OTHERWISE LISTED UNDER THIS CHEMICAL] | T*0.1 |

| **HALOSULFURON-METHYL**<br>HALOSULFURON-METHYL | |
|---|---|
| EDIBLE OFFAL (MAMMALIAN) | 0.2 |
| MEAT (MAMMALIAN) | *0.01 |
| MILKS | *0.01 |

| **IMAZAPIC**<br>SUM OF IMAZAPIC AND ITS HYDROXYMETHYL DERIVATIVE | |
|---|---|
| EGGS | *0.01 |
| POULTRY, EDIBLE OFFAL OF | *0.01 |
| POULTRY MEAT | *0.01 |

| **METALAXYL**<br>METALAXYL | |
|---|---|
| EDIBLE OFFAL (MAMMALIAN) | *0.05 |
| MILKS | *0.01 |

| **METHOMYL**<br>SUM OF METHOMYL AND METHYL HYDROXYTHIOACETIMIDATE ('METHOMYL OXIME'), EXPRESSED AS METHOMYL<br>*SEE ALSO* THIODICARB | |
|---|---|
| CORIANDER (LEAVES, STEM, ROOTS) | T10 |
| HERBS | T10 |

| **NORFLURAZON**<br>NORFLURAZON | |
|---|---|
| ASPARAGUS | 0.05 |

| **PIRIMICARB**<br>SUM OF PIRIMICARB, DIMETHYL-PIRIMICARB AND N-FORMYL-(METHYLAMINO) ANALOGUE (DIMETHYLFORMAMIDIO-PIRIMICARB), EXPRESSED AS PIRIMICARB | |
|---|---|
| LEAFY VEGETABLES | T5 |

| **PROCYMIDONE**<br>PROCYMIDONE | |
|---|---|
| ADZUKI BEAN (DRY) | T0.2 |
| BEANS [EXCEPT BROAD BEAN AND SOYA BEAN] | T10 |
| EDIBLE OFFAL (MAMMALIAN) | T0.05 |
| EGGS | T*0.01 |
| GARLIC | T5 |
| LUPIN (DRY) | T*0.01 |
| MEAT (MAMMALIAN) (IN THE FAT) | T0.2 |
| MILKS | T0.02 |
| ONION, BULB | T0.2 |
| POME FRUITS | T1 |
| POTATO | T0.1 |
| POULTRY, EDIBLE OFFAL OF | T*0.01 |
| POULTRY MEAT (IN THE FAT) | T0.1 |
| SNOW PEAS | T5 |
| STONE FRUITS | T10 |

| **PROPACHLOR**<br>PROPACHLOR | |
|---|---|
| RADISH | *0.02 |
| SWEDE | *0.02 |

| **SETHOXYDIM**<br>SUM OF SETHOXYDIM AND METABOLITES CONTAINING THE 5-(2-ETHYLTHIOPROPYL)CYCLOHEXENE-3-ONE AND 5-HYDROXYCYCLOHEXENE-3-ONE MOIETIES AND THEIR SULFOXIDES AND SULFONES, EXPRESSED AS SETHOXYDIM | |
|---|---|
| BRASSICA (COLE OR CABBAGE) VEGETABLES, HEAD CABBAGES, FLOWERHEAD BRASSICAS | 0.5 |
| LETTUCE, HEAD | 0.2 |
| LETTUCE, LEAF | 0.2 |
| PEANUT | 3 |

| **TRICHLORFON**<br>TRICHLORFON | |
|---|---|
| MILKS | *0.05 |

| **TRIFLOXYSTROBIN**<br>SUM OF TRIFLOXYSTROBIN AND ITS ACID METABOLITE ((E,E)-METHOXYIMINO-[2-[1-(3-TRIFLUOROMETHYLPHENYL)-ETHYLIDENEAMINOOXYMETHYL]PHENYL] ACETIC ACID), EXPRESSED AS TRIFLOXYSTROBIN EQUIVALENTS | |
|---|---|
| STRAWBERRY | 2 |